

Geochemical and source characteristics of the Tintagel Volcanic Formation

B. RICE-BIRCHALL and P.A. FLOYD



Rice-Birchall, B. and Floyd, P.A. 1988. Geochemical and source characteristics of the Tintagel Volcanic Formation. *Proceedings of the Ussher Society*, 7, 52-55.

The Lower Carboniferous metavolcanic greenschists of the Tintagel Volcanic Formation (TVF) are confirmed as a single cogenetic suite of fractionated alkali basalt lavas and volcanoclastics. The volcanics are highly altered and absolute element abundances have been diluted in highly carbonated samples, although ratios of stable incompatible elements remain constant. The TVF displays the following chemical features: uniform incompatible element ratios ($Zr/Nb \sim 6$; $La/Nb \sim 0.8$; $La/Ta \sim 10$) and light REE (chondrite-normalized) enrichment with relatively uniform $(La/Yb)_N \sim 6-8$; in this respect they are comparable with Upper Devonian basalts from N Cornwall and Devon. Relative to MORB the TVF rocks exhibit progressive incompatible element enrichment patterns with a positive Nb-Ta anomaly that are characteristic of an intraplate environment and an enriched OIB-type mantle source. There is no chemical evidence for crustal contamination.

B. Rice-Birchall and P.A. Floyd, Department of Geology, University of Keele, Staffordshire, ST5 5BG.

Introduction

The Tintagel Volcanic Formation (TVF) is a series of highly foliated meta-lavas and volcanoclastics (Dearman *et al.* 1970; Freshney *et al.* 1972), exposed on the north Cornish coast between Trebarwith Strand and Boscastle. The local stratigraphy within the Tredorn Nappe (Stewart 1981) comprises a Dinantian succession of the Barras Nose Formation, the Tintagel Volcanic Formation and the Trambley Cove Formation, all of which are tectonically overlain by light grey Upper Devonian slates. The sequence has a polyphase deformation history (Sanderson 1979; Selwood and Thomas 1986) and was metamorphosed in the lower greenschist facies (Robinson and Read 1981) as indicated by the characteristic assemblage of chlorite-muscovite-calcite-epidote-biotite.

Recently, Robinson and Sexton (1987), using statistical techniques on a small chemical data set collected from the TVF, demonstrated that most major and some trace elements were mobile during post-magmatic alteration and metamorphism. However, using the least mobile trace elements they showed that the TVF represented a single, fractionated basic suite of alkali basalt lineage. This primary feature of the TVF is confirmed by our data collected from throughout the outcrop.

Objectives

Some 59 samples of metavolcanic greenschists and associated metasediments were collected from the TVF outcrop at Trebarwith Strand, Barras Nose and Gullastem and analysed for major and trace elements by XRF Spectrometry (at University of Keele), REE (by ICP Source Spectrometry at R.H.B.N. College, University of London) and Hf, Sc, U, Th and Ta (by INAA at Universities Research Reactor, Risley). A representative selection of analysed TVF samples is shown in Table 1.

We wished to determine (a) whether the TVF represented a single chemical suite (cf. Robinson and Sexton 1987) throughout its entire outcrop, (b) if the metavolcanic greenschists contained a chemically recognizable contaminant from either the associated metasediments or continental crust, and (c) the nature of the source region and subsequent eruptive environment relative to Upper Devonian metabasalts from south-west England.

Lithological variation

Within the metavolcanics two main lithotypes can be recognised: (a) homogeneous, fine-grained, well foliated, completely recrystallised greenschists, and (b) heterogeneous, often coarser-grained, layered greenschists with variably sized lenses and phacoids of basaltic lava in a recrystallized foliated matrix. Petrographically, the less-deformed lenses exhibit relict primary

volcanic textures including glassy matrix, quenched crystals and phenocrysts. Preliminary examination indicates that both plagioclase phyric basalts and minor fine-grained dolerites constituted the original volcanics. Although they have an overall basic composition and could have originally represented a series of pillow lavas and sills, they could equally have been accumulations of pillow breccias and tuffaceous volcanoclastics with basaltic clasts.

Geochemical features

Chemical fractionation

Previous work (Robinson and Sexton 1987) showed that the TVF is a cogenetic alkali basalt suite with chemical variation governed by olivine-plagioclase fractionation. These results are confirmed by our data which show linear covariance between pairs of stable incompatible elements indicative of a single chemical series with alkaline characteristics. The comagmatic nature of the TVF is also supported by REE patterns that are typical for rocks related via crystal fractionation. Samples from all the main outcrops show parallel, light REE-enriched (chondrite-normalized) patterns indicative of either plagioclase (petrographically observed) or olivine fractionation. However there is no evidence to support the suggestion that the TVF is composed of less-evolved material at the top of the volcanic sequence (Robinson and Sexton 1987). The lower absolute abundances recorded are the result of differential dilution by secondary calcite in this part of the sequence. In particular, samples with high calcite contents (> 15 modal%), have their trace element concentrations considerably reduced. However, the ratio between any two immobile incompatible elements remains constant suggesting the ratio has not been changed by alteration. In this context element ratios are generally a more reliable geochemical parameter than the absolute abundances. In particular the TVF is characterized by relatively constant immobile incompatible element ratios such as, $Zr/Nb \sim 6$, $La/Nb \sim 0.8$ and $(La/Yb)_N \sim 6-8$, displaying a moderate light-to-heavy REE fractionation.

Incompatible element patterns and mantle sources

The top diagram of Fig. 1 is a "spidergram" showing four representative TVF samples normalized to MORB values (after Saunders and Tarney 1984) for a range of incompatible elements, together with Ni and Cr. A number of features are exhibited:

(a) The samples have closely parallel patterns with most elements being progressively enriched relative to MORB. The general parallelism implies that the samples from the three main localities are genetically related by fractional crystallization. The large negative Sr anomaly is most probably an alteration feature, although together with the small negative Eu anomaly could be indicative of plagioclase fractionation, which is confirmed by the

Table 1: Representative chemical data for metavolcanic greenschists from the Tintagel Volcanic Formation. Major elements in wt%, trace elements in ppm.

Sample number	TV-1	TV-2	TV-9	TV-10	TV-15	TV-16	TV-35	TV-36
Locality code*	BN	BN	G	G	TS	TS	TS	TS
SiO ₂	43.8	46	41.5	44.3	38.8	37.9	43.9	40.9
TiO ₂	3.32	2.68	2.01	2.45	1.76	1.64	2.41	2.09
Al ₂ O ₃	21.4	17	14.8	15.2	13.6	12.4	14.5	14
Fe ₂ O _{3t}	15.1	13.3	12.2	11.6	10.9	10.8	9.52	12.5
MnO	0.14	0.12	0.16	0.13	0.11	0.12	0.11	0.13
MgO	4.7	6.47	8.58	6.93	10.3	11.4	1.95	3.17
CaO	1.81	4.29	8.7	7.83	9.4	9.86	10.9	11.1
Na ₂ O	0.2	0.38	1.3	3.59	1.76	1.03	4.84	2.71
K ₂ O	4.16	2	1.73	0.71	1.42	1.4	2.05	2.3
P ₂ O ₅	0.31	0.47	0.31	0.53	0.26	0.16	0.6	0.33
LOI	5.29	6.94	8.01	7.59	11.8	12.7	9.28	11
Total	100	99.6	99.3	101	100	99.4	100	100
Ba	31	30	307	29	37	72	450	303
Cr	173	124	150	256	438	408	120	49
Cu	12	17	95	44	53	82	23	29
Ga	27	20	17	17	19	16	19	21
Nb	42	36	28	37	22	23	28	29
Ni	73	76	106	144	254	244	51	46
Pb	31	25	17	15	10	11	12	15
Rb	132	60	36	11	27	30	56	58
Sr	85	83	468	306	158	163	267	292
V	197	159	224	214	202	187	268	183
Y	40	42	30	36	26	24	39	33
Zn	163	157	111	106	92	98	72	101
Zr	330	274	161	224	129	125	180	171
Cs	10.2	7.3	5.6	3	6.6	9.6	2.42	3.45
Hf	8.2	6.5	4	5.5	3.5	3.2	4.8	4.2
Sc	29.2	23.6	30.3	23.6	27.5	25.1	23.2	20.1
Ta	2.85	2.53	1.66	2.48	1.4	1.37	1.9	1.83
Th	2.9	2.49	2.16	3.09	2.02	1.9	2.37	1.96
U	0.79	0.67	0.67	0.95	0.71	0.62	1.09	0.78
La	24.7	28.8	15.8	26	14	12.5	18.2	24.5
Ce	66.1	68.9	38.2	59.8	33.3	30.1	46	52.4
Pr	8.2	8.54	4.75	7.3	4.2	3.71	6.27	6.46
Nd	38.6	40.2	22.8	33.6	19.9	17.8	30.5	29.9
Sm	8.49	8.59	4.74	6.71	4.16	3.7	6.56	5.96
Eu	2.63	2.67	1.63	2.12	1.33	1.23	2.05	1.7
Gd	8.68	9.34	5.42	7.08	4.73	4.3	7.31	6.41
Dy	7.42	7.59	4.9	5.89	4.16	3.79	6.11	5.18
Ho	1.47	1.53	1.03	1.2	0.86	0.8	1.25	1.06
Er	3.9	4.11	2.86	3.31	2.41	2.23	3.44	2.95
Yb	2.94	3.25	2.3	2.65	1.91	1.82	2.7	2.28
Lu	0.37	0.45	0.31	0.36	0.27	0.25	0.35	0.3

* BN = Barras Nose, G = Gullastem, TS = Trebarwith Strand

presence of plagioclase phenocrysts in many lava lenses. Variation in normalized Ni and Cr also suggest mafic fractionation between the illustrated samples, with TV-18 being the most primitive. The greater degree of scatter shown by K, Ba and Rb is due to mobility during alteration.

(b) The relatively enriched "humped" pattern with a pronounced Nb and Ta positive anomaly is characteristic of within plate settings (Pearce 1983) and in particular ocean island basalts (OIB) and their mantle source (Wood *et al.* 1979; Bougault *et al.* 1980; Tarney *et al.* 1980; Hofmann 1986). In the lower diagram TV-18 is compared with two primitive alkali basalts from within plate oceanic settings which precludes contamination by continental crust. Although the absolute abundances of elements vary, the shapes of the three patterns are quite similar and suggest that the TVF volcanics were derived from an enriched OIB-type mantle source, clearly distinct from a MORB-type source.




Figure 1. MORB-normalized multi-element diagram of representative basic volcanics from different sectors of the Tintagel Volcanic Formation, N. Cornwall. Lower diagram compares a typical Tintagel volcanic (TV-18) with intraplate alkali basalts from Annobon Island, Gulf of Guinea (Liottard *et al.* 1982) and Suiko seamount, Emperor/Hawaiian chain (Cambon *et al.* 1980). MORB normalizing factors from Saunders and Tamey (1984).

An enriched source for TVF relative to primitive mantle is indicated by Nb/U (~50) and La/Ta (~10) ratios that are remarkably similar to OIB sources (~47 and ~9 respectively, Bougault *et al.* 1980; Hofmann 1986). Hofmann (1986) has suggested that a Nb-Ta-enriched source could be generated with the retention of these elements in refractory Ti oxides during continental crust formation at island arcs. These enriched portions of mantle are then recycled as OIB "plums" (Morris and Hart 1983) and form the source for oceanic island basalts.

Crustal contamination

The TVF and the associated pelitic metasediments form two discrete chemical groups indicating that there is no mixing between them. Also, the contents of P, Ba and Th in the volcanics do not reach the high levels expected if contaminated with pelitic sediments. Contamination by continental crust is often associated with a negative Nb-Ta anomaly (Thompson *et al.* 1983) and consequently an increase in La/Ta and La/Nb coupled with a decrease in Nb/U, none of which are observed in the TVF.

At this stage and without the benefit of isotopic data contamination with continental crust and country rock is thought to have played a negligible part in the evolution of the TVF.

Eruptive setting

The secondary calcite effect can have important consequences for the discrimination of tectonic environment if only the Ti-Y-Zr diagram (Pearce and Cann 1973) is used. As seen in Fig. 2 the calcite-poor samples plot in the within-plate field (cf. Robinson and Sexton 1987), whereas the calcite-rich samples are pulled towards the Y apex and apparently suggest a different tectonic environment. Although excess calcite proportionally dilutes the




Figure 2. Variable distribution of Tintagel volcanics in the tectonic discrimination diagram of Pearce and Cann (1973) depending on their secondary carbonate content. Fields are: 1 = intraplate basalts; 2 = predominantly arc tholeiites; 3 = predominantly mid-ocean ridge basalts; 4 = predominantly arc calc-alkali basalts.

absolute abundances of all covariant trace elements, in extreme cases Y can be mobilized via CO₂ complexing and slightly enriched during secondary carbonation.

Using only calcite-poor samples, the diagrams in Fig. 3 both demonstrate the within-plate (and alkaline) character of the TVF. The high La/Y and Nb/Y ratios are typical of alkaline provinces in continental environments and suggest the Lower Carboniferous basin was ensialic (cf. Leeder 1982). With the exception of some metabasalts from S. Cornwall, a within-plate tectonic setting is characteristic of Hercynian volcanics from Cornubia (Floyd 1982, 1983).

Comparison with Upper Devonian basalts

Floyd (1984) demonstrated that the incompatible element chemistry of Cornubian basaltic volcanics varied with tectonic province. As exhibited in Fig. 4 Upper Devonian and S. Cornish melange basalts can be discriminated in terms of Zr and Nb. The Lower Carboniferous TVF from N. Cornwall plots with their Upper Devonian analogues from the same area and suggests a similar mantle source was being tapped throughout Upper Palaeozoic volcanic events.

Conclusions

1. The TVF is a single, cogenetic, fractionated alkaline basaltic suite of lavas and/or basic volcanoclastics.
2. They exhibit (MORB-normalized) progressive incompatible element enrichment patterns characterized by a positive Nb-Ta anomaly and a negative Sr anomaly. Their chemical features indicate they were derived from an enriched OIB-type mantle source and erupted in a within-plate setting.
3. There is no trace element evidence for crustal contamination or pelitic sediment mixing in the volcanics.
4. The TVF rocks have an incompatible element chemistry similar to Upper Devonian basalts from N. Cornwall and Devon.

Acknowledgements. The University of Keele Research Committee is acknowledged for providing field work expenses. D. Emley and M. Aiken are thanked for technical help in the production of the XRF data and Dr. N. Walsh (RHBNC) for the REE data.

References

- Bougault, H., Joron, J.-L. and Treuil, M. 1980. The primordial chondritic nature and large scale heterogeneities in the mantle: evidence from high and low partition coefficient elements in oceanic basalts. *Philosophical Transactions of the Royal Society*, A297, 203-213.
- Cambon, P., Joron, J.-L., Bougault, H. and Treuil, M. 1980. Leg 55, Emperor seamounts: trace elements in transitional tholeiites, alkali basalts and hawaiites - mantle homogeneity or heterogeneity arid magmatic processes. In: Jackson et al (eds) Initial Reports of the *Deep Sea Drilling Project*, 55, U.S. Government Printing Office, Washington, 585-597.
- Dearman, W.R., Freshney, E.C., King, A.F., Williams, M. and McKeown, M.C. 1970. The north coast of Cornwall from Bude to Tintagel. *Geologists' Association Guides*, No. 10, 1-42.
- Floyd, P.A. 1982. Chemical variation in Hercynian basalts relative to plate tectonics. *Journal of the Geological Society*, London, 139, 505-520.
- Floyd, P.A. 1983. Composition and petrogenesis of the Lizard complex pre-orogenic basaltic rocks in southwest England. In: Hancock, P.L. (ed) *The variscan fold belt in the British Isles*. Adam Hilger Ltd., Bristol, 130-152.
- Floyd, P.A. 1984. Geochemical characteristics and comparison of the basic rocks of the Lizard complex and the basaltic lavas within the Hercynian troughs of SW England. *Journal of the Geological Society*, London, 141, 61-70.
- Freshney, E.C., McKeown, M.C. and Williams, M. 1972. Geology of the coast between Tintagel and Bude. *Memoir of the Geological Survey of Great Britain*.
- Hofmann, A.W. 1986. Nb in Hawaiian magmas: constraints on source composition and evolution. *Chemical Geology*, 57, 17-19.
- Leeder, M. 1982. Upper Palaeozoic basins of the British Isles - Caledonide inheritance versus Hercynian plate margin processes. *Journal of the Geological Society*, London, 139, 479-491.
- Liotard, J.M., Dupuy, C., Dostal, J. and Cornen, G. 1982. Geochemistry of the volcanic island of Annobon, Gulf of Guinea. *Chemical Geology*, 35, 115-128.
- Morris, J.D. and Hart, S.R. 1983. Isotopic and incompatible element constraints on the genesis of island arc volcanics from Cold Bay and Amak Islands, Aleutians, and implications for mantle structure. *Geochimica et Cosmochimica Acta*, 47, 2015-2030.
- Pearce, J.A. 1983. Role of the sub-continental lithosphere in magma genesis at active continental margins. In: Hawkesworth, C.J. and Norry, M.J. (eds) *Continental basalts and mantle xenoliths*. Shiva, Cheshire, 230-249.




Figure 3. Chemical discrimination of the Tintagel basic volcanics eruptive setting as a within-plate alkaline province. Immobile element ratio diagram based on data from the literature (Floyd, unpublished) and only includes tholeiitic basalts for the oceanic island environment; oceanic alkali basalts overlap with the continental field that includes both magma types. Hf-Ta-Th diagram from Wood (1980).




Figure 4. Comparison of Lower Carboniferous Tintagel basic volcanics with Upper Devonian basalts from Cornwall and Devon, as well as S. Cornish melange metabasalts and Start Point greenschists. Comparative data from Floyd (1983) and unpublished.

Pearce, J.A. and Cann, J.R. 1973. Tectonic setting of basic volcanic rocks determined using trace element analysis. *Earth and Planetary Science Letters*, 19, 290-300.

Robinson, D. and Read, D. 1981. Metamorphism and mineral chemistry of greenschists from Trebarwith Strand, Cornwall. *Proceedings of the Ussher Society*, 5, 132-138.

Robinson, D. and Sexton, D. 1987. Geochemistry of the Tintagel Volcanic Formation. *Proceedings of the Ussher Society*, 6, 523-528.

Sanderson, D.J. 1979. The transition from upright to recumbent folds in the Variscan fold belt of southwest England: a model based on the kinematics of simple shear. *Journal of Structural Geology*, 1, 171-180.

Saunders, A.D. and Tarney, J. 1984. Geochemical characteristics of basaltic volcanism within back-arc basins. In: Kokelaar, B.P. and Howells, M.F. (eds) *Marginal basin geology. Geological Society Special Publication No. 16*, 59-76.

Selwood, E.B. and Thomas, J.M. 1986. Upper Palaeozoic successions and nappe structures in north Cornwall. *Journal of the Geological Society*, 143, 75-82.

Stewart, I.J. 1981. The Trekellend Thrust. *Proceedings of the Ussher Society*, 5, 163-167.

Tarney, J., Wood, D.A., Saunders, A.D., Cann, J.R. and Varet, J. 1980. Nature of mantle heterogeneity in the North Atlantic: evidence from deep sea drilling. *Philosophical Transactions of the Royal Society*, London, A297, 179-202.

Thompson, R.N., Morrison, M.A., Dickin, A.P. and Hendry, G.L. 1983. Continental flood basalts... Arachnids rule OK? In: Hawkesworth, C.J. and Norry, M.J. (eds) *Continental basalts and mantle xenoliths*. Shiva, Cheshire.

Wood, D.A. 1980. The application of a Th-Hf-Ta diagram to problems of tectomagmatic classification and to establishing the nature of crustal contamination of basaltic lavas of the British Tertiary volcanic province. *Earth and Planetary Science Letters*, 50, 11-30.

Wood, D.A., Joron, J.-L., Treuil, M., Norry, M. and Tarney, J. 1979. Elemental and Sr isotopic variations in basic lavas from Iceland and the surrounding ocean floor. *Contributions to Mineralogy and Petrology*, 70, 319-339.